## **CLINITEST Rapid COVID-19 Antigen Self-Test**

**Quick Reference Instructions** 

This test is free and distributed by the U.S. Health Resources & Services Administration (HRSA) through NEMS to ensure and expand free and accessible testing to the community. It is not authorized for re-sale.


\*This test is authorized for self-testing use by individuals aged 12 and older (or adult-collected samples from individuals ages 2-13).

The US Food and Drug Administration has granted a shelf-life extension for the CLINITEST® **Rapid COVID-19 Antigen Self-Test** from 11 months to 24 months from the date of manufacture.

> Please scan the QR CODE for updated information.




STFP 1



Before starting, wash your hands. Place tube in the holder and remove the seal. Make sure the tube is standing straight.

5 circles



5 circles

Carefully insert swab tip about 1/2 to 3/4 inch deep into nostril and complete at least 5 circles in EACH nostril.

STEP 3



Insert swab tip into the tube and mix vigorously at least 6 times on the bottom and side of the tube.

STEP 4




Place tube in the tube holder. Start timer for 1 minute. After 1 minute, squeeze swab tip several times and remove from the tube. Insert tube tip into the top of the tube.


STEP 5 Result window Sample well

Squeeze 4 drops of solution into the sample well and start timer for 15 minutes.

## **READ TEST RESULT AFTER 15 MINUTES**



COVID-19 POSITIVE: Two pink/red lines indicate a positive result. Immediately notify your health care provider about your positive result to discuss COVID-19 treatment options. Note that oral antiviral treatment for COVID-19 must be started within 5 days of symptom onset.



COVID-19 NEGATIVE: COVID-19 viral proteins were not detected. Consider retesting in 1-2 days if your symptoms started within 3 days ago. NEMS NORTH EAST

**INVALID:** This test result is invalid. Perform a new test.

北醫療 a california health.t center